



U.S. ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE

USAMRICD-TR-99-02

Measure of O⁶-Alkylguanine-DNA
Alkyltransferase Activity in Normal
Human Epidermal Keratinocytes in
Culture and Effects of *Bis*-(2-chloroethyl)
sulfide on the Activity

Ellen M. Kurt
Vincent M. O'Donnell
Charise C. Davenport
Agnes Gaweska

19990813 015

May 1999

Approved for public release; distribution is unlimited

U.S. Army Medical Research
Institute of Chemical Defense
Aberdeen Proving Ground, MD 21010-5400

DISPOSITION INSTRUCTIONS:

Destroy this report when no longer needed. Do not return to the originator.

DISCLAIMERS:

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

The use of trade names does not constitute an official endorsement or approval of the use of such commercial hardware or software. This document may not be cited for purposes of advertisement.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE May 1999		3. REPORT TYPE AND DATES COVERED January 1997 to September 1998
4. TITLE AND SUBTITLE Measure of O ⁶ -Alkylguanine-DNA Alkyltransferase Activity in Normal Human Epidermal Keratinocytes in Culture and Effects of Bis-(2-chloroethyl)sulfide on the Activity.			5. FUNDING NUMBERS TC1	
6. AUTHOR(S) Kurt, E.M., O'Donnell, V.M., Davenport, C.C., and Gaweska, A.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-DA 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-RC 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400			10. SPONSORING/MONITORING AGENCY REPORT NUMBER USAMRICD-TR-99-02	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) O ⁶ -Alkylguanine-DNA alkyltransferase (AGT) is a DNA repair protein that removes alkyl adducts from O ⁶ -alkylguanine in DNA. AGT may be important in DNA repair following injury induced by bis-(2-chloroethyl)sulfide (sulfur mustard, HD), since O ⁶ -alkylguanine is one of the HD alkylation products. One model for HD-induced injury uses normal human epidermal keratinocytes (NHEK) in culture. In this study we measured the levels of AGT in NHEK grown to either 60- 80% or to 100% confluency and then studied the effects of a one-hour exposure to 50, 100, and 300 μ M HD on AGT activity. Mean AGT activity in NHEK grown to 60-80% and 100% confluency was 470 \pm 404 and 518 \pm 737 fmol/mg protein, respectively. In general, AGT activity appeared to increase after exposure to 100 μ M HD and decrease with increased confluency and after exposure to 50 and 300 μ M HD. However, a two-way analysis of variance for cell confluency and HD concentration showed no significant differences between cell confluencies or the different HD concentrations.				
14. SUBJECT TERMS Sulfur mustard, DNA damage, DNA repair, AGT, normal human epidermal keratinocytes, O ⁶ -methylguanine-DNA methyltransferase			15. NUMBER OF PAGES 12	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UNLIMITED	

ACKNOWLEDGEMENTS

The authors wish to acknowledge Ms. Theresa Nipwoda for growing the NHEK, Mr. Offie Clark for help with HD exposures, Dr. William Smith for useful discussions and help with HD exposures, and Ms. Robyn Lee for help with the statistical analysis.

INTRODUCTION

Sulfur mustard (HD) is a potent alkylating agent that has a long history of use as a chemical warfare agent. Despite decades of medical research, the mechanism by which HD induces vesication is not known, and no effective antidotes are currently available. The fact that DNA is a major cellular target for HD alkylation has been well documented (Fox and Scott, 1980; Rozmiarek et al., 1973). HD alkylates DNA to form ethylthioethyl adducts to include monofunctional adducts at the O⁶ position of guanine (Papirmeister et al., 1991).

O⁶-Alkylguanine-DNA alkyltransferase (AGT) is a repair protein that acts by transferring the alkyl group from O⁶-alkylguanine to a cysteine residue in its center. The reaction is stoichiometric: each molecule of AGT removes one alkyl group, which becomes covalently bound to the AGT cysteine residue (Brent, 1985; Lindahl et al., 1988; Yarosh, 1985; Pegg and Dolan 1987). The alkylated transferase molecule is then catabolized. The cell's ability to perform this type of repair depends strictly on the level of AGT (Yarosh, 1985; Pegg and Dolan, 1987). The level of AGT varies greatly among species, and different tissues contain different levels of alkyltransferase activity (Gerson et al., 1986).

Previous studies have established that normal human epidermal keratinocytes (NHEK) in culture can serve as useful models for biochemical investigations into the toxicity of HD (Smith et al., 1990). Here we report the AGT activity in cultured NHEK grown to either 60-80% or to 100% confluency, and the effects of a one-hour exposure to 50, 100, and 300 μ M HD on the AGT activity.

METHODS

Sulfur mustard. Bis-(2-chloroethyl)sulfide (5 μ l in 10 ml of Keratinocyte Growth Medium [KGM]) was obtained from the U.S. Army Edgewood Research, Development and Engineering Center (Aberdeen Proving Ground, MD).

Cell culture. Primary cultures of adult NHEK (strain 4075) were obtained from Clonetics Corporation (San Diego, CA). Tertiary cultures of NHEK were passed into 150 cm² tissue culture flasks and grown until either 60-80% or 100% confluency. At this time 50, 100, or 300 μ M HD was added to half the flasks, while the other half of the flasks served as the controls. Seven samples were exposed to 50 μ M HD, seven samples to 100 μ M HD, and seven samples to 300 μ M HD. The flasks were left at room temperature for one hour and then harvested by trypsinization according to the procedure described by Boyce and Ham (1985). The harvested NHEK were collected in a conical tube and placed in a styrofoam box containing dry ice until AGT analysis. The cell counts, determined using trypan blue staining, ranged from 2.0x10⁷ to 4.6x10⁷ cells/culture/sample.

Assay of AGT activity. The activity of AGT in NHEK was measured in the laboratory of Dr. Eileen Dolan at the Division of Hematology-Oncology, the University of Chicago Medical Center (Chicago, Illinois) as previously described (Dolan et al., 1990). Briefly, the NHEK cells were incubated with [³H]methylated DNA, which was prepared by allowing [³H]N-methylnitrosourea to react with calf thymus DNA. The DNA was precipitated then acid hydrolyzed, and the modified bases were separated by reverse-phase HPLC. Protein was determined by the method of Bradford (1976), and the results are expressed as fmol of O⁶-methylguanine released from the DNA substrate per mg protein. Each sample was run in

duplicate. The results are reported as the mean \pm the standard deviation. The AGT activity in NHEK exposed for one hour to 50, 100, or 300 μ M HD is reported as percent change of control where % change = [(control-HD)/control] x 100.

RESULTS

Table I shows the mean AGT activity measured for the cultured NHEK. Mean AGT activity measured in 12 samples of NHEK grown to 60-80% confluency was 470 ± 404 fmol/mg protein. Mean AGT activity measured in 6 samples of NHEK grown to 100% confluency was 518 ± 737 fmol/mg protein. A two-way analysis of variance showed no significant differences between confluency levels. Because there were no significant differences, all AGT activities were grouped together and the mean calculated. The mean AGT activity for the 18 samples was 485 ± 515 fmol/mg protein, which is not significantly different from zero.

Table II shows the percent changes of the control of AGT activity in NHEK, at 60-100% confluency, for the HD-exposed samples. The percent changes of the control of the AGT activity for the 50, 100, and 300 μ M HD exposed NHEK were $-18 \pm 167\%$, $+52 \pm 98.1\%$, and $-72 \pm 86.9\%$, respectively. A two-way analysis of variance showed no significant differences among the HD exposure concentrations (as compared with the controls). Because there were no significant differences among HD exposure concentrations, the percent changes for all three different concentrations were grouped together and averaged. The mean percent change following exposure of the NHEK to HD is $10.06 \pm 88.9\%$, which is not significantly different from zero.

DISCUSSION

This is the first report of AGT activity in cultured NHEK as well as in NHEK exposed to HD. The AGT activity in 60-80% confluent control NHEK (470 ± 404 fmol/mg protein) is higher than that reported for other human tissues, such as brain (67 ± 40 fmol/mg protein), lung (86 fmol/mg protein), myeloid precursors (126 ± 59 fmol/mg protein), colon (138 ± 78 fmol/mg protein), and small intestine (242 ± 168 fmol/mg protein). It is, however, comparable to that reported for T-lymphocytes (359 ± 72 fmol/mg protein) and liver (485 ± 126 fmol/mg protein) (Gerson et al., 1986).

In general, AGT activity decreased with an increase in confluency. The trends for change in AGT activity in NHEK following a one-hour HD exposure were for an increase in activity after 100 μ M HD and a decrease in activity after 50 and 300 μ M HD. None of these trends was supported by statistical analyses, which showed no significant differences in AGT activities between NHEK of 60-80% and 100% confluency and no significant differences in AGT activity in unexposed NHEK versus NHEK exposed to 50, 100, or 300 μ M HD. While the statistical analyses show that AGT activity is not a function of cell confluency and that a one-hour exposure to 50, 100, and 300 μ M HD does not affect AGT activity in NHEK in culture, the general trends do show some correlation indicating that a greater sample population may provide statistical support for the observed trends.

Table 1. Mean O⁶-Alkylguanine-DNA Alkyltransferase (AGT) Activity
in Normal Human Epidermal Keratinocytes in Culture

Number of NHEK Samples	% Confluency	AGT Activity (fmol/mg protein)
12	60 - 80	470 ± 404
6	100	518 ± 737
18 ^a	60 - 100	485 ± 515

^aThe mean AGT activity for the 18 NHEK samples.

Table II. Percent Changes of Control of O⁶-Alkylguanine-DNA Alkyltransferase Activity in Normal Human Epidermal Keratinocytes, at 60-100% Confluency, Exposed to Different Sulfur Mustard (HD) Concentrations

Number of Determinations	HD Concentrations (μ M)	Percent Change ^b
7	50	-18 \pm 167
7	100	+ 52 \pm 98.1
7	300	-72 \pm 86.9

^aPercent change = [(control - HD exposed)/control] x 100

REFERENCES

- Boyce, S. and Ham, R. (1985) Procedure No. 41406. *J. Tissue Culture Methods*. **9**, 85-93.
- Brent, T. (1985). Isolation and purification of O⁶-alkylguanine-DNA alkyltransferase from human leukemic cells. Prevention of chloroethylnitrosourea-induced cross-links by purified enzyme. *Pharmacol. Ther.* **31**, 121-140.
- Bradford, M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principles of protein-dye binding. *Anal. Biochem.* **72**, 248-254.
- Dolan, M., Moschel, R., and Pegg, A. (1990). Depletion of mammalian O⁶-alkylguanine-DNA alkyltransferase activity by O⁶-benzylguanine provides a means to evaluate the role of this protein in protection against carcinogenic and therapeutic alkylating agents. *Proc. Natl. Acad. Sci. USA* **87**, 5368-5372.
- Fox, M. and Scott, D. (1980). The genetic toxicology of nitrogen and sulphur mustard. *Mutation Research* **75**, 131-168.
- Gerson, S., Trey, J., Miller, K., and Berger, N. (1986). Comparison of O⁶-alkylguanine-DNA alkyltransferase activity based on cellular DNA content in human, rat and mouse tissues. *Carcinogenesis* **7**, 745-749.
- Lindahl, T., Sedgwich, B., Sekiguchi, M. and Nakabeppu, Y. (1988). Regulation and expression of the adaptive response to alkylating agents. *Annu. Rev. Biochem.* **57**, 133-157.
- Papirmeister, B., Feister, A., Robinson, S., and Ford, R. (1991) *Medical Defense Against Mustard Gas*, pp. 104-107. CRC Press: Boca Raton.
- Pegg, A. and Dolan, M. (1987). Properties and assay of mammalian O⁶-alkylguanine-DNA alkyltransferase. *Pharmacol. Ther.* **34**, 167-179.
- Rozmiarek, J., Cappizzi, R., Papirmeister, B., Furman, W., and Smith, W. (1973). Mutagenic activity in somatic and germ cells following chronic inhalation of sulfur mustard. *Mutat. Res.* **21**, 13-14.
- Smith, W., Gross, C., Chan, P., and Meier, H. (1990). The use of human epidermal keratinocytes in culture as a model for studying the biochemical mechanisms of sulfur mustard toxicity. *Cell Biol. Toxicol.* **6**, 285-290.
- Yarosh, D. (1985). The role of O⁶-methylguanine-DNA methyltransferase in cell survival, mutagenesis and carcinogenesis. *Mutat. Res.* **145**, 1-16.

DISTRIBUTION LIST

Addresses

Copies

DEFENSE TECHNICAL INFORMATION
CENTER
ATTN DTIC OCP
8725 JOHN J KINGMAN RD STE 0944
FT BELVOIR VA 22060-6218

2

DIRECTOR
ARMED FORCES MEDICAL
INTELLIGENCE CENTER
1607 PORTER STREET
FORT DETRICK MD 21702-5004

1

COMMANDER
US ARMY MEDICAL RESEARCH AND
MATERIEL COMMAND
ATTN MCMR PLD
504 SCOTT ST
FORT DETRICK MD 21702-5012

2

COMMANDER
US ARMY INSTITUTE OF DENTAL RESEARCH
BUILDING 40
WASHINGTON DC 20307-5300

1

HQDA
OFFICE OF THE SURGEON GENERAL
5109 LEESBURG PIKE SUITE 691
FALLS CHURCH VA 22041-3258

1

COMMANDER
US ARMY INSTITUTE OF SURGICAL
RESEARCH
BUILDING 2653
FORT SAM HOUSTON TX 78234-6200

1

DIRECTOR
WALTER REED ARMY INSTITUTE OF
RESEARCH
BUILDING 40
WASHINGTON DC 20307-5100

1

COMMANDER
USAMEDD CENTER & SCHOOL
ATTN MCCS FC
FORT SAM HOUSTON TX 78234-6100

1

COMMANDER
US ARMY AEROMEDICAL RESEARCH
LABORATORY
ATTN SCIENTIFIC INFORMATION CENTER
PO BOX 577
FORT RUCKER AL 36362-5000

1

COMMANDER
USAMEDD CENTER & SCHOOL
ATTN MCCS FCD
FORT SAM HOUSTON TX 78234-6100

1

COMMANDER
US ARMY MEDICAL RESEARCH INSTITUTE
OF INFECTIOUS DISEASES
1425 PORTER ST
FORT DETRICK MD 21702-5011

1

DIRECTOR
ENVIRONMENTAL AND LIFE SCIENCES
OFFICE OF THE DEPUTY DIRECTOR FOR
RESEARCH AND ENGINEERING
ROOM 3D129
WASHINGTON DC 20301-2300

1

COMMANDER
US ARMY RESEARCH INSTITUTE
OF ENVIRONMENTAL MEDICINE
ATTN MCMR UE ZS (MS SAFRAN)
BUILDING 42
NATICK MA 01760-5007

1

COMMANDER
US ARMY TRAINING AND DOCTRINE
COMMAND
ATTN ATMD
FORT MONROE VA 23651

1

COMMANDANT
US ARMY CHEMICAL SCHOOL
ATTN ATZN CM C
FORT MCCLELLAN AL 36205

1

COMMANDER
US ARMY NUCLEAR AND CHEMICAL AGENCY
7500 BACKLICK ROAD
BUILDING 2073
SPRINGFIELD VA 22150-3198

1

COMMANDER 1
US ARMY MEDICAL MATERIEL
DEVELOPMENT ACTIVITY
ATTN MCMR UMD
622 NEIMAN ST
FORT DETRICK MD 21702-5009

EXECUTIVE OFFICER 1
NAVAL MEDICAL RESEARCH INSTITUTE
NAVAL MEDICINE COMMAND
NATIONAL CAPITAL REGION
BETHESDA MD 20814

DEPARTMENT OF THE NAVY 1
NAVAL POSTGRADUATE SCHOOL
DUDLEY KNOX LIBRARY
411 DYER ROAD
ROOM 110
MONTEREY CA 93943-5101

USAF ARMSTRONG LABORATORY/CFTO 1
SUSTAINED OPERATIONS BRANCH
BROOKS AFB TX 78235-5000

DEPARTMENT OF HEALTH AND HUMAN 1
SERVICES
NATIONAL INSTITUTES OF HEALTH
THE NATIONAL LIBRARY OF MEDICINE
SERIAL RECORDS SECTION
8600 ROCKVILLE PIKE
BETHESDA MD 20894

STEMSON LIBRARY 1
ACADEMY OF HEALTH SCIENCES
BUILDING 2840 RM 106
FORT SAM HOUSTON TX 78234-6100

US ARMY RESEARCH OFFICE 1
ATTN CHEMICAL AND BIOLOGICAL
SCIENCES DIVISION
PO BOX 12211
RESEARCH TRIANGLE PARK NC 27709-2211

AFOSR/NL 1
BUILDING RM A217
BOLLING AFB DC 20332

COMMANDER 1
US ARMY CHEMICAL BIOLOGICAL
DEFENSE AGENCY
ATTN AMSCB CI
ABERDEEN PROVING GROUND MD
21010-5423

LTC RICHARD R. STOTTS 1
BATTELLE MEMORIAL INSTITUTE JM 3
505 KING AVENUE
COLUMBUS OH 43201-2695

COMMANDER 23
US ARMY MEDICAL RESEARCH
INSTITUTE OF CHEMICAL DEFENSE
3100 RICKETTS POINT ROAD
ATTN MCMR UV ZA
MCMR UV ZB
MCMR UV ZS
MCMR UV RC (5 copies)
MCMR UV R (11 copies)
MCMR UV AI W
MCMR UV D
MCMR UV P
MCMR UV C
ABERDEEN PROVING GROUND MD
21010-5425